

DRINKING WATER SOURCE PROTECTION

ACT FOR CLEAN WATER

Thames-Sydenham and Region Source Protection Committee
Upper Thames River Source Protection Area

Amended Proposed
Assessment Report

Revised - August 12, 2011

APPROVED

2.0 Watershed Characterization



Ontario

Made possible through the support
of the Government of Ontario





Thames - Sydenham and Region
Upper Thames River Conservation Authority
1424 Clarke Road, London, ON, N5V 5B9

July 4, 2011

Dear reader

Re: **Upper Thames River Amended Proposed Assessment Report posted for comments**

The Thames-Sydenham and Region Source Protection Committee has posted the enclosed report for review and comment by stakeholders. This report includes updates and amendments to the Proposed Assessment Report for the Upper Thames River Source Protection Authority. Comments received through the first posting (draft proposed report) were considered by the Source Protection Committee and the report has been revised. Comments received in response to the second posting were submitted with the Proposed Assessment Report to the Ministry of the Environment (MOE).

Based on directions from the Director of Source Protection Programs Branch, MOE, updates and amendments were made to the Proposed Assessment Report to include further technical work conducted. These include the Tier 2 water budget drought scenario analysis, the delineation, vulnerability assessment and threats assessment of WHPA-E, and threats assessment of sewer line threats. The updates and amendments made are summarized in Appendix 4. This version of the Assessment Report once approved will be used by the Source Protection Committee for the development of the Source Protection Plan which is to be submitted to the MOE in August 2012.

The Amended Proposed Assessment Report for the Upper Thames River Source Protection Area represents a significant milestone in the Source Protection Committee's progress in the completion of the first Source Protection Plans for the Thames-Sydenham and Region. The Source Protection Committee realizes that this report is a "living document" which may be updated as more information becomes available in the future.

We hope that you have had an opportunity to attend previous open houses that were held throughout the region, and plan to attend the open houses to be held in July 2011. More information on this assessment report and the past and current consultation is available on the web site given at the bottom of this letter.

Yours truly,
THAMES-SYDENHAM and REGION

Robert Bedgood, Chair
Source Protection Committee

Lower Thames Valley Conservation Authority
100 Thames Street, Chatham, Ontario,
N7L 2Y8

phone 519-354-7310, fax 519-352-3435

St. Clair Region Conservation Authority
205 Mill Pond Cres., Strathroy, Ontario,
N7G 3P9

phone 519-245-3710, fax. 519-245-3348

Upper Thames River Conservation Authority
1424 Clarke Road, London, ON
N5V 5B9

phone 519-451-2800, fax 519-451-1188

Upper Thames River Source Protection Area Assessment Report

Table of Contents

Appendices.....	iv
List of Figures.....	iv
List of Maps.....	iv
List of Tables.....	vii
1.0 Introduction and Background.....	1-1
1.1 Document Overview.....	1-2
1.2 Clean Water Act Rules and Regulations.....	1-5
1.2.1 Regulations.....	1-5
1.2.2 Technical Rules.....	1-6
1.2.3 Local Guidance Documents.....	1-6
1.2.4 Tables of Drinking Water Threats.....	1-6
1.2.5 Mapping Symbology.....	1-7
1.2.6 Source Protection Plan.....	1-7
1.3 Source Protection Committee.....	1-8
1.4 Role of the Conservation Authorities.....	1-10
1.5 Terms of Reference.....	1-11
1.6 Thames-Sydenham and Region Source Protection Region.....	1-12
1.6.1 Upper Thames River Source Protection Area.....	1-12
1.7 Technical Studies.....	1-13
1.8 Consultation.....	1-14
1.9 Schedule.....	1-16
1.10 Local Acceptance, Approvals and Next Steps.....	1-17
1.10.1 Engaging First Nations.....	1-19
1.10.2 Amendments to the Assessment Report.....	1-20
2.0 Watershed Characterization.....	2-1
2.1 Watershed Characterization Report.....	2-1
2.2 Data Sources.....	2-2
2.3 Components of the Watershed Characterization Report.....	2-4
2.3.1 Watersheds and Subwatersheds.....	2-4
2.3.2 Physical Geography.....	2-4
2.3.3 Human Geography.....	2-14
2.3.4 Water Quality.....	2-18
2.3.5 Water Quantity.....	2-22
2.3.6 Drinking Water Systems.....	2-22
2.4 Data Gaps.....	2-28
3.0 Water Budget and Water Quantity Stress Assessment.....	3-1
3.1 What is a Water Budget?.....	3-2
3.2 Components of the Water Budget.....	3-3
3.2.1 Precipitation.....	3-3
3.2.2 Evapotranspiration.....	3-3
3.2.3 Surface Runoff.....	3-4
3.2.4 Recharge.....	3-4
3.2.5 Water Use (Demand).....	3-4
3.2.6 Water Budget Summary.....	3-9

Upper Thames River Source Protection Area Assessment Report

3.3	Phases of Water Budget Work	3-10
3.3.1	Conceptual Water Budget	3-10
3.3.2	Tier 1 Water Budget	3-11
3.3.3	Tier 2 Water Budget	3-12
3.3.4	Tier 3 Water Budget	3-13
3.3.5	Peer Review of the Water Budget	3-13
3.4	Water Quantity Stress Assessment.....	3-14
3.4.1	Uncertainty in the Stress Assessment.....	3-21
3.5	Significant Groundwater Recharge Areas	3-22
3.6	Data Gaps and Next Steps.....	3-24
4.0	Vulnerability Assessment.....	4-1
4.1	Peer Review of Vulnerability Assessment.....	4-2
4.2	Intake Protection Zones	4-3
4.3	Wellhead Protection Areas	4-3
4.3.1	Technical Studies	4-4
4.3.2	WHPA-A	4-7
4.3.3	WHPA-B, WHPA-C and WHPA-D	4-7
4.3.4	WHPA-E and WHPA-F	4-7
4.3.5	Vulnerability Assessment of the WHPA.....	4-23
4.3.6	Uncertainty in WHPA Vulnerability	4-47
4.4	Highly Vulnerable Aquifers	4-48
4.4.1	Uncertainty of HVA	4-52
4.5	Significant Groundwater Recharge Areas	4-52
4.5.1	Uncertainty of SGRA	4-54
4.6	Data Gaps and Next Steps.....	4-54
5.0	Issues Evaluation	5-1
5.1	What is a Drinking Water Quality Issue?	5-1
5.2	Impact of Identifying an Issue.....	5-6
5.3	Issue Evaluation Methodology	5-6
5.4	Issues Evaluation Technical Studies	5-9
5.5	Identified Issues	5-10
5.6	Work Plan.....	5-14
5.7	Data Gaps	5-14
6.0	Conditions Assessment	6-1
6.1	Conditions Assessment Methodology	6-3
6.1.1	Situations Where Conditions May Exist.....	6-3
6.1.2	Information Used to Identify Conditions.....	6-4
6.1.3	Risk Assessment Methodology for Conditions	6-5
6.2	Conditions Assessment Findings	6-7
6.3	Data Gaps and Next Steps for Conditions	6-8
7.0	Threats and Risk Assessment – Water Quality	7-1
7.1	Drinking Water Quality Threat Identification and Risk Assessment Methodology.....	7-3
7.1.1	Prescribed Drinking Water Threats	7-4
7.1.2	Other Activities	7-14
7.1.3	Threats Arising from Conditions	7-14
7.1.4	Threats Arising from Issues.....	7-15

Upper Thames River Source Protection Area Assessment Report

7.1.5	Local Guidance and Technical Studies	7-16
7.2	Drinking Water Quality Threats and Risk Assessment.....	7-17
7.2.1	Threats Identified through Mapping of Impervious Surfaces, Managed Lands and Livestock Density	7-18
7.2.2	Number of Locations of Significant Threats.....	7-18
7.2.3	Threats in Birr Wellhead Protection Areas	7-22
7.2.4	Threats in Dorchester Wellhead Protection Areas	7-22
7.2.5	Threats in Kilworth-Komoka Wellhead Protection Areas – WELLS DECOMMISSIONED	7-23
7.2.6	Threats in City of London Wellhead Protection Areas.....	7-24
7.2.7	Threats in Melrose Wellhead Protection Areas	7-25
7.2.8	Threats in Thorndale Wellhead Protection Areas.....	7-25
7.2.9	Threats in Beachville Wellhead Protection Areas.....	7-26
7.2.10	Threats in Embro Wellhead Protection Areas	7-27
7.2.11	Threats in Hickson Wellhead Protection Areas	7-28
7.2.12	Threats in Ingersoll Wellhead Protection Areas	7-28
7.2.13	Threats in Innerkip Wellhead Protection Areas	7-29
7.2.14	Threats in Lakeside Wellhead Protection Areas.....	7-30
7.2.15	Threats in Mount Elgin Wellhead Protection Areas	7-31
7.2.16	Threats in Tavistock Wellhead Protection Areas.....	7-31
7.2.17	Threats in Thamesford Wellhead Protection Areas.....	7-32
7.2.18	Threats in Woodstock Wellhead Protection Areas	7-33
7.2.19	Threats in Mitchell Wellhead Protection Areas.....	7-34
7.2.20	Threats in Sebringville Wellhead Protection Areas	7-35
7.2.21	Threats in Shakespeare Wellhead Protection Areas.....	7-36
7.2.22	Threats in St. Marys Wellhead Protection Areas.....	7-37
7.2.23	Threats in St. Pauls Wellhead Protection Areas.....	7-38
7.2.24	Threats in Stratford Wellhead Protection Areas	7-38
7.2.25	Threats in HVA and SGRA.....	7-39
7.3	Tier 2 Risk Assessment.....	7-40
7.4	Data Gaps	7-40
8.0	Great Lakes.....	8-1
8.1	Impact of Considering Great Lakes.....	8-2
8.2	Great Lakes Agreements	8-3
8.2.1	Great Lakes Water Quality Agreement.....	8-3
8.2.2	The Canada-Ontario Agreement Respecting the Great Lakes Basin Ecosystem ..	8-5
8.2.3	The Great Lakes Charter and the Great Lakes-St. Lawrence River Basin Sustainable Water Resources Agreement.....	8-6
8.3	Proposed Working Groups	8-7
8.4	Next Steps for Great Lakes	8-8
9.0	Data Gaps and Next Steps.....	9-1
9.1	Data Gaps	9-1
9.2	Next Steps.....	9-4

Upper Thames River Source Protection Area Assessment Report

Appendices

- Appendix 1 – Maps (bound separately)
- Appendix 2 – Section Summaries (bound separately)
- Appendix 3 – System Summaries (bound separately)
- Appendix 4 – Assessment Report Consultation
- Appendix 5 – Watershed Characterization Summary (bound separately)
- Appendix 6 – Conceptual Water Budget (bound separately)
- Appendix 7 – Assessment Report Checklist
- Appendix 8 – Issues Evaluation Methodology
- Appendix 9 – Issues Evaluation Flagged Parameters
- Appendix 10 – Threats and Risk Assessment
- Appendix 11 – Glossary of Terms and Acronyms
- Appendix 12 – References
- Appendix 13 – Uncertainty Analysis

List of Figures

- Figure 1-1 Source Protection planning schedule overview1-17
- Figure 5-1 Thames-Sydenham and Region Issues Evaluation Methodology5-8

List of Maps

- 1-1 Thames-Sydenham and Region Source Protection Region
- 1-2 Upper Thames River Source Protection Area
- 1-3 Municipal Drinking Water Systems
- 1-4 Areas of Settlement
- 2-1 Population Density
- 2-2 Federal Lands
- 3-1 Thames-Sydenham & Region Tier 1 Water Budget Subwatersheds
- 3-2 Average Precipitation Distribution
- 3-3 Average Evapotranspiration Distribution
- 3-4 Mean Annual Recharge
- 3-5 Surface Water Potential for Stress
- 3-6 Groundwater Potential for Stress
- 4-1 Vulnerability Overview of Wellhead Protection Area (WHPA)
 - 4-1-1 Birr Wellhead Protection Area (WHPA)
 - 4-1-2 Dorchester Wellhead Protection Area (WHPA)
 - 4-1-2a Dorchester Wellhead Protection Area (WHPA-E)
 - 4-1-3 Kilworth-Komoka Wellhead Protection Area (WHPA) - DECOMMISSIONED
 - 4-1-4 London – Fanshawe Wellhead Protection Area (WHPA)
 - 4-1-4a London – Fanshawe Wellhead Protection Area (WHPA-E)
 - 4-1-5 London – Hyde Park Wellhead Protection Area (WHPA)
 - 4-1-6 Melrose Wellhead Protection Area (WHPA)
 - 4-1-7 Thorndale Wellhead Protection Area (WHPA)

Upper Thames River Source Protection Area Assessment Report

- 4-1-8 Beachville Wellhead Protection Area (WHPA)
- 4-1-9 Embro Wellhead Protection Area (WHPA)
- 4-1-10 Hickson Wellhead Protection Area (WHPA)
- 4-1-11 Ingersoll Wellhead Protection Area (WHPA)
- 4-1-12 Innerkip Wellhead Protection Area (WHPA)
- 4-1-13 Lakeside Wellhead Protection Area (WHPA)
- 4-1-14 Mount Elgin Wellhead Protection Area (WHPA)
- 4-1-15 Tavistock Wellhead Protection Area (WHPA)
- 4-1-16 Thamesford Wellhead Protection Area (WHPA)
- 4-1-16a Thamesford Wellhead Protection Area (WHPA-E)
- 4-1-17 Woodstock Wellhead Protection Area (WHPA)
- 4-1-17a Woodstock Wellhead Protection Area (WHPA-E)
- 4-1-18 Mitchell Wellhead Protection Area (WHPA)
- 4-1-19 Sebringville Wellhead Protection Area (WHPA)
- 4-1-20 Shakespeare Wellhead Protection Area (WHPA)
- 4-1-21 St. Marys Wellhead Protection Area (WHPA)
- 4-1-21a St. Marys Wellhead Protection Area (WHPA-E)
- 4-1-22 St. Pauls Wellhead Protection Area (WHPA)
- 4-1-23 Stratford Wellhead Protection Area (WHPA)
- 4-2-1 Significant Groundwater Recharge Areas
- 4-2-2 Significant Groundwater Recharge Areas (SGRA) Vulnerability
- 4-3-1 Aquifer Vulnerability
- 4-3-2 Highly Vulnerable Aquifers (HVA)
- 7-1-1 Birr Drinking Water Threats Impervious Surface Areas
- 7-1-2 Dorchester Drinking Water Threats Impervious Surface Areas
- 7-1-3 Kilworth-Komoka Drinking Water Threats Impervious Surface Areas - DECOMMISSIONED
- 7-1-4 London – Fanshawe Drinking Water Threats Impervious Surface Areas
- 7-1-5 London – Hyde Park Drinking Water Threats Impervious Surface Areas
- 7-1-6 Melrose Drinking Water Threats Impervious Surface Areas
- 7-1-7 Thorndale Drinking Water Threats Impervious Surface Areas
- 7-1-8 Beachville Drinking Water Threats Impervious Surface Areas
- 7-1-9 Embro Drinking Water Threats Impervious Surface Areas
- 7-1-10 Hickson Drinking Water Threats Impervious Surface Areas
- 7-1-11 Ingersoll Drinking Water Threats Impervious Surface Areas
- 7-1-12 Innerkip Drinking Water Threats Impervious Surface Areas
- 7-1-13 Lakeside Drinking Water Threats Impervious Surface Areas
- 7-1-14 Mount Elgin Drinking Water Threats Impervious Surface Areas
- 7-1-15 Tavistock Drinking Water Threats Impervious Surface Areas
- 7-1-16 Thamesford Drinking Water Threats Impervious Surface Areas
- 7-1-17 Woodstock Drinking Water Threats Impervious Surface Areas
- 7-1-18 Mitchell Drinking Water Threats Impervious Surface Areas
- 7-1-19 Sebringville Drinking Water Threats Impervious Surface Areas
- 7-1-20 Shakespeare Drinking Water Threats Impervious Surface Areas
- 7-1-21 St. Marys Drinking Water Threats Impervious Surface Areas
- 7-1-22 St. Pauls Drinking Water Threats Impervious Surface Areas
- 7-1-23 Stratford Drinking Water Threats Impervious Surface Areas
- 7-1-24 Impervious Surface Area in Highly Vulnerable Aquifers (HVA)
- 7-1-25 Impervious Surface Area in Significant Groundwater Recharge Areas (SGRA)
- 7-2-1 Birr Drinking Water Threats Managed Lands and Livestock Density
- 7-2-2 Dorchester Drinking Water Threats Managed Lands and Livestock Density
- 7-2-3 Kilworth-Komoka Drinking Water Threats Managed Lands and Livestock Density - DECOMMISSIONED
- 7-2-4 London – Fanshawe Drinking Water Threats Managed Lands and Livestock Density

Upper Thames River Source Protection Area Assessment Report

- 7-2-5 London – Hyde Park Drinking Water Threats Managed Lands and Livestock Density
- 7-2-6 Melrose Drinking Water Threats Managed Lands and Livestock Density
- 7-2-7 Thorndale Drinking Water Threats Managed Lands and Livestock Density
- 7-2-8 Beachville Drinking Water Threats Managed Lands and Livestock Density
- 7-2-9 Embro Drinking Water Threats Managed Lands and Livestock Density
- 7-2-10 Hickson Drinking Water Threats Managed Lands and Livestock Density
- 7-2-11 Ingersoll Drinking Water Threats Managed Lands and Livestock Density
- 7-2-12 Innerkip Drinking Water Threats Managed Lands and Livestock Density
- 7-2-13 Lakeside Drinking Water Threats Managed Lands and Livestock Density
- 7-2-14 Mount Elgin Drinking Water Threats Managed Lands and Livestock Density
- 7-2-15 Tavistock Drinking Water Threats Managed Lands and Livestock Density
- 7-2-16 Thamesford Drinking Water Threats Managed Lands and Livestock Density
- 7-2-17 Woodstock Drinking Water Threats Managed Lands and Livestock Density
- 7-2-18 Mitchell Drinking Water Threats Managed Lands and Livestock Density
- 7-2-19 Sebringville Drinking Water Threats Managed Lands and Livestock Density
- 7-2-20 Shakespeare Drinking Water Threats Managed Lands and Livestock Density
- 7-2-21 St. Marys Drinking Water Threats Managed Lands and Livestock Density
- 7-2-22 St. Pauls Drinking Water Threats Managed Lands and Livestock Density
- 7-2-23 Stratford Drinking Water Threats Managed Lands and Livestock Density
- 7-2-24 Percent Managed Lands in Highly Vulnerable Aquifers (HVA)
- 7-2-25 Livestock Density in Highly Vulnerable Aquifers (HVA)
- 7-2-26 Percent Managed Lands in Significant Groundwater Recharge Area (SGRA) with Vulnerability Score = 6
- 7-2-27 Livestock Density in Significant Groundwater Recharge Area (SGRA) with Vulnerability Score = 6
- 7-3-1 Birr Areas Where Activities Related to Pathogens, Chemicals or DNAPL are or would be Drinking Water Threats
- 7-3-2 Dorchester Areas Where Activities Related to Pathogens, Chemicals or DNAPL are or would be Drinking Water Threats
- 7-3-2a Dorchester (WHPA-E) Areas Where Activities Related to Pathogens, Chemicals or DNAPL are or would be Drinking Water Threats
- 7-3-3 Kilworth-Komoka Areas Where Activities Related to Pathogens, Chemicals or DNAPL are or would be Drinking Water Threats - DECOMMISSIONED
- 7-3-4 London – Fanshawe Areas Where Activities Related to Pathogens, Chemicals or DNAPL are or would be Drinking Water Threats
- 7-3-4a London – Fanshawe (WHPA-E) Areas Where Activities Related to Pathogens, Chemicals or DNAPL are or would be Drinking Water Threats
- 7-3-5 London – Hyde Park Areas Where Activities Related to Pathogens, Chemicals or DNAPL are or would be Drinking Water Threats
- 7-3-6 Melrose Areas Where Activities Related to Pathogens, Chemicals or DNAPL are or would be Drinking Water Threats
- 7-3-7 Thorndale Areas Where Activities Related to Pathogens, Chemicals or DNAPL are or would be Drinking Water Threats
- 7-3-8 Beachville Areas Where Activities Related to Pathogens, Chemicals or DNAPL are or would be Drinking Water Threats
- 7-3-9 Embro Areas Where Activities Related to Pathogens, Chemicals or DNAPL are or would be Drinking Water Threats
- 7-3-10 Hickson Areas Where Activities Related to Pathogens, Chemicals or DNAPL are or would be Drinking Water Threats
- 7-3-11 Ingersoll Areas Where Activities Related to Pathogens, Chemicals or DNAPL are or would be Drinking Water Threats
- 7-3-12 Innerkip Areas Where Activities Related to Pathogens, Chemicals or DNAPL are or would be Drinking Water Threats

Upper Thames River Source Protection Area Assessment Report

- 7-3-13 Lakeside Areas Where Activities Related to Pathogens, Chemicals or DNAPL are or would be Drinking Water Threats
- 7-3-14 Mount Elgin Areas Where Activities Related to Pathogens, Chemicals or DNAPL are or would be Drinking Water Threats
- 7-3-15 Tavistock Areas Where Activities Related to Pathogens, Chemicals or DNAPL are or would be Drinking Water Threats
- 7-3-16 Thamesford Areas Where Activities Related to Pathogens, Chemicals or DNAPL are or would be Drinking Water Threats
- 7-3-16a Thamesford (WHPA-E) Areas Where Activities Related to Pathogens, Chemicals or DNAPL are or would be Drinking Water Threats
- 7-3-17 Woodstock Areas Where Activities Related to Pathogens, Chemicals or DNAPL are or would be Drinking Water Threats
- 7-3-17a Woodstock (WHPA-E) Areas Where Activities Related to Pathogens, Chemicals or DNAPL are or would be Drinking Water Threats
- 7-3-18 Mitchell Areas Where Activities Related to Pathogens, Chemicals or DNAPL are or would be Drinking Water Threats
- 7-3-19 Sebringville Areas Where Activities Related to Pathogens, Chemicals or DNAPL are or would be Drinking Water Threats
- 7-3-20 Shakespeare Areas Where Activities Related to Pathogens, Chemicals or DNAPL are or would be Drinking Water Threats
- 7-3-21 St. Marys Areas Where Activities Related to Pathogens, Chemicals or DNAPL are or would be Drinking Water Threats
- 7-3-21a St. Marys (WHPA-E) Areas Where Activities Related to Pathogens, Chemicals or DNAPL are or would be Drinking Water Threats
- 7-3-22 St. Pauls Areas Where Activities Related to Pathogens, Chemicals or DNAPL are or would be Drinking Water Threats
- 7-3-23 Stratford Areas Where Activities Related to Pathogens, Chemicals or DNAPL are or would be Drinking Water Threats

List of Tables

Table 1-1 SPC members and representation.....	1-9
Table 1-2 SPC Mission Statement and Guiding Principles	1-10
Table 1-3 Municipalities in the UTRSPA	1-12
Table 1-4 Summary of planned UTRSPA Assessment Report Consultation	1-16
Table 1-5 First Nations of the TSR SPR	1-19
Table 2-1 Watershed Characterization Report Data Sources	2-2
Table 2-2 Habitat and Temperature Assessments in the UTRSPA	2-9
Table 2-3 Population Density of Municipalities in the Upper Thames River SPA Watershed ..	2-14
Table 2-4 Municipal Drinking Water Systems Serving the Upper Thames River Source Protection Area	2-24
Table 2-5 Watershed Characterization Data Gaps relevant to the Upper Thames River Source Protection Area	2-29
Table 3-1 Groundwater use in the UTRSPA (m ³ /day).....	3-7
Table 3-2 Surface Water use in the UTRSPA (m ³ /day)†	3-8
Table 3-3 Tier 2 water budget summary (m ³ /day).....	3-10
Table 3-4 Potential for stress based on <i>percent water demand</i> under current and future municipal water demand	3-17

Upper Thames River Source Protection Area Assessment Report

Table 3-5 Surface water potential for stress based on Tier 1 stress assessment (Month of August).....	3-18
Table 3-6 Groundwater potential for stress (Average Annual Conditions).....	3-18
Table 3-7 Groundwater potential for stress (Maximum Monthly Conditions).....	3-19
Table 3-8 Data gaps related to Water Budget and Water Quantity Stress Assessment.....	3-24
Table 4-1 Summary of Technical Reports for UTR Municipal Systems.....	4-5
Table 4-2 Thames-Sydenham and Region GUDI wells.....	4-8
Table 4-3 WHPA vulnerability scoring (Technical Rules Table 2 a and b).....	4-41
Table 4-4 List of Groundwater Wells and Vulnerability Scores for WHPA-A, B, C, D.....	4-41
Table 4-5 List of GUDI Wells and Vulnerability Scores for WHPA-E.....	4-43
Table 4-6 Vulnerability Assessment Data Gaps Relevant to the Upper Thames River SPA...4-54	
Table 5-1 Schedule 1 Parameters (from O. Reg. 169/03 of the Safe Drinking Water Act, 2002) and their Treated Drinking Water Quality Standards.....	5-2
Table 5-2 Schedule 2 Parameters (O. Reg. 169/03 of the Safe Drinking Water Act, 2002) and their Treated Drinking Water Quality Standards.....	5-3
Table 5-3 Schedule 3 Parameters (O. Reg. 169/03 of the Safe Drinking Water Act, 2002) and their Treated Drinking Water Quality Standards.....	5-4
Table 5-4 Table 4 Parameters (from the Technical Support Document for the Ontario Drinking Water Standards, Objectives and Guidelines, MOE 2006) with their Treated Drinking Water Aesthetic Objectives and Operational Guidelines.....	5-5
Table 5-5 Technical Studies on Drinking Water Quality Issues Evaluation.....	5-9
Table 5-6 Drinking Water Quality Issues Identified in the Upper Thames River Source Protection Area.....	5-11
Table 5-7 Determination of Sources of an Issue.....	5-15
Table 6-1 Technical Studies on Drinking Water Threats and Risk Assessment.....	6-2
Table 6-2 Threat Level Determination for Conditions.....	6-6
Table 7-1 Technical Studies on Drinking Water Threats and Risk Assessment.....	7-2
Table 7-2 Activities Prescribed as Drinking Water Threats.....	7-4
Table 7-3 Threat Level Determination.....	7-6
Table 7-4 Chemical Hazard Scorings for Various Combinations of Percentage of Managed Lands and Livestock Densities.....	7-12
Table 7-5 Number of Locations of Significant Drinking Water Threats in Middlesex County and City of London.....	7-19
Table 7-6 Number of Locations of Significant Drinking Water Threats in Oxford County.....	7-20
Table 7-7 Number of Locations of Significant Drinking Water Threats in Perth County, City of Stratford and Town of St. Marys.....	7-21
Table 7-8 Number of Locations of Significant Drinking Water Threats in HVA and SGRA.....	7-21
Table 7-9 Significant Threats in the Birr WHPA.....	7-22
Table 7-10 Significant Threats in the Dorchester WHPA.....	7-23
Table 7-11 Significant Threats in the Kilworth-Komoka WHPA - WELLS DECOMMISSIONED 7-23	
Table 7-12 Significant Threats in the City of London-Fanshawe WHPA.....	7-24
Table 7-13 Significant Threats in the City of London-Hyde Park WHPA.....	7-24
Table 7-14 Significant Threats in the Melrose WHPA.....	7-25
Table 7-15 Significant Threats in the Thorndale WHPA.....	7-26
Table 7-16 Significant Threats in the Beachville WHPA.....	7-27
Table 7-17 Significant Threats in the Embro WHPA.....	7-27
Table 7-18 Significant Threats in the Hickson WHPA.....	7-28

Upper Thames River Source Protection Area Assessment Report

Table 7-19 Significant Threats in the Ingersoll WHPA	7-29
Table 7-20 Significant Threats in the Innerkip WHPA.....	7-30
Table 7-21 Significant Threats in the Lakeside WHPA	7-30
Table 7-22 Significant Threats in the Mount Elgin WHPA.....	7-31
Table 7-23 Significant Threats in the Tavistock WHPA	7-32
Table 7-24 Significant Threats in the Thamesford WHPA	7-33
Table 7-25 Significant Threats in the Woodstock WHPA (Urban well system)	7-34
Table 7-26 Significant Threats in the Woodstock WHPA (Rural well system)	7-34
Table 7-27 Significant Threats in the Mitchell WHPA	7-35
Table 7-28 Significant Threats in the Sebringville WHPA	7-36
Table 7-29 Significant Threats in the Shakespeare WHPA	7-36
Table 7-30 Significant Threats in the St. Marys WHPA	7-37
Table 7-31 Significant Threats in the St. Pauls WHPA	7-38
Table 7-32 Significant Threats in the Stratford WHPA.....	7-39
Table 7-33 Levels of Threats Related to Pathogens, Chemicals and DNAPLs in HVAs and SGRAs	7-40
Table 9-1 Work Plan to fill Data and Analysis Gaps	9-2

2.0 Watershed Characterization

Justice O'Connor recommended that watershed-based *Source Protection Plans* be developed. The recommendations were part of the inquiry which investigated the May 2000 bacterial contamination of the Town of Walkerton's water supply. Compiling a summary of information pertinent to drinking water sources is one of the first steps in developing a *Source Protection Plan*.

Under the Clean Water Act (2006), the Assessment Report must identify all subwatersheds in the source protection area and characterize the water quality and quantity across the watershed. The Regulations and *rules* under the Clean Water Act (2006) require that the physical and human geography also be characterized. This information is contained in a watershed characterization report.

2.1 Watershed Characterization Report

The Watershed Characterization Report for the Thames Watershed and Region, completed in 2008, is based on information available at the time. Updated characterization information is included in other sections of the Assessment Report. Some of the water budget related mapping products are available in the Conceptual Water Budget, which is included as an appendix to the Assessment Report.

The Watershed Characterization Report summarizes information on the physical, social and economic characteristics of the Thames Watershed & Region. It reviews surface water and groundwater quality, and summarizes known issues and concerns pertaining to drinking water sources. A series of maps help to illustrate the information presented in the report. Each of the components of the watershed characterization report is described in the sections that follow.

Upper Thames River Source Protection Area Assessment Report

The summary of the Watershed Characterization Report for the Thames Watershed and Region is included in Appendix 5 in the Lower Thames Valley and Upper Thames River Source Protection Area Assessment Reports; complete with all maps. The entire Watershed Characterization Report is available on compact disk (CD).

2.2 Data Sources

A wide range of data sources have been used as resources to prepare the Watershed Characterization Report and the accompanying maps. Data used to characterize the Thames watershed is provided in Table 2-1 below.

Table 2-1 Watershed Characterization Report Data Sources

Component	Data Source
Bedrock Geology	Waterloo Hydrogeologic. 2005. Six Conservation Authorities FEFLOW Groundwater Model: Conceptual Model Report.
Surficial Geology	Waterloo Hydrogeologic. 2005. Southwestern Region Edge-Matching Study. Surficial Geology of Southern Ontario. Ontario Geological Survey Miscellaneous Release –Data 128.
Physiography	Chapman, L.J. and D.F. Putnam. 1984. The Physiography of Southern Ontario, 3rd edition.
Soils Information	Ontario Ministry of Agriculture and Food and Agriculture Canada, Soils Ontario Version 1.0. Ontario Soils Surveys
Groundwater Hydrogeology	Waterloo Hydrogeologic. 2005. Six Conservation Authorities FEFLOW Groundwater Model: Conceptual Model Report. Waterloo Hydrogeologic. 2005. Southwestern Region Edge-Matching Study. Municipal Groundwater Studies. MOE.
Surface Water Hydrology	Ontario Ministry of the Environment and Ontario Ministry of Natural Resources. 1975. Thames River Basin Water Management Study. Stream Gauge Data. Ontario Ministry of Agriculture and Food and Agriculture. Municipal Drain Classification (Fisheries and Oceans Canada project) data. UTRCA. 1991. Dam Inventory and Reservoir Assessment.
Naturally Vegetated Areas	Ministry of Natural Resources Aerial Mapping 2001 and 2003.
Aquatic Ecology	Species at Risk Recovery Plan. Fisheries and Oceans Canada. Ontario Ministry of Natural Resources. Royal Ontario Museum. Ontario Ministry of the Environment. COA and COA partners - Thames River Habitat Assessment and Monitoring Program. Aquatic Species at Risk in the Thames River Watershed, Ontario. Cudmore, B., C. A. MacKinnon and S. E. Madzia. Dec. 2004. Canadian Manuscript Report of

Upper Thames River Source Protection Area Assessment Report

Table 2-1 Watershed Characterization Report Data Sources

Component	Data Source
	<p>Fisheries and Aquatic Sciences. 2707.</p> <p>Thames River Recovery Team. 2004. Recovery strategy for the Thames River Aquatic Ecosystem: 2005-2010. December 2004 Draft. 145 pp. Natural Heritage Information Centre.</p> <p>COSEWIC assessment and status report on the Rainbow mussel <i>Villosa iris</i> in Canada. COSEWIC. 2006. Committee on the Status of Endangered Wildlife in Canada. Ottawa. Vii + 38 pp. (www.sararegistry.gc.ca/status/status_e.cfm).</p> <p>Spiny Softshell and Queen Snake Research and Recovery Along the Thames River Watershed. Gillingwater, S.D. 2009. Report submitted to the Ontario Ministry of Natural Resources.</p> <p>National Recovery Strategy for the Queen Snake (<i>Regina septemvittata</i>) in Canada. Gillingwater, S.D. 2008. Prepared for the Queen Snake Recovery Team. Draft. 38 pp.</p> <p>Rare Reptile Research of the Thames River Watershed. Gillingwater, S.D. and T.J. Piraino. 2002. Report submitted to the Ontario Ministry of Natural Resources, Aylmer District.</p> <p>Freshwater mussel communities of the Thames River, Ontario: 2004-2005. Morris, T.J. and A. Edwards. 2007. Can. Manuscr. Rpt. Fish. Aquat. Sci. 2810: v + 30 pp.</p> <p>Queen Snakes (<i>Regina septemvittata</i>) and Spiny Softshell Turtles (<i>Apalone spinifera spinifera</i>) Along the Upper Thames River Watershed. Piraino, T.J. and S.D. Gillingwater. 2004. Report submitted to the Upper Thames River Conservation Authority.</p>
Human Characterization	<p>Statistics Canada. Censuses of Population, 1901-2001 and 1996-2006.</p> <p>Indian and Northern Affairs Canada website: http://ainc-inac.gc.ca</p> <p>Ontario Ministry of Finance Ontario Population Projections, 2006-2031.</p> <p>Municipality Official Plans.</p> <p>Ministry of Environment. June 1991. Waste Disposal Site Inventory.</p> <p>Census Canada.</p> <p>Ontario Ministry of Agriculture and Food and Agriculture.</p>
Drinking Water Sources	<p>Ministry of Environment Permit To Take Water (PTTW) database.</p> <p>Municipal Groundwater Studies. MOE.</p>
Water Quality	<p>Provincial Water Quality Monitoring Network.</p> <p>Provincial Groundwater Monitoring Network.</p> <p>Drinking Water Surveillance Program.</p> <p>Drinking Water Information System.</p> <p>Annual Drinking Water System Reports.</p> <p>Ministry of Environment Inspection reports.</p> <p>Water treatment plant laboratory data.</p> <p>Ambient Groundwater Chemistry Study of the Thames River and St. Clair Region Watersheds. Waterloo Hydrologic Incorporated, 2008.</p>

2.3 Components of the Watershed Characterization Report

2.3.1 Watersheds and Subwatersheds

The source protection area (*SPA*) watershed boundary within the source protection region (SPR), as well as the subwatersheds within the *SPA*, are identified and described. The Thames watershed and region is comprised of the Lower Thames Valley Source Protection Area (*LTVSPA*) and the Upper Thames River Source Protection Area (*UTRSPA*). Map 1-1 in Appendix 1 illustrates the Thames-Sydenham and Region boundary and the Source Protection Area watershed boundaries within the Region.

The Upper Thames River Source Protection Area includes all areas draining into the Thames River above the community of Delaware. This area covers large parts of Oxford, Perth and Middlesex Counties, including most of the City of London. Very small portions of Huron and Elgin Counties also drain into the upper Thames River. The *UTRSPA* covers approximately 3,423 square kilometres with a total watershed population (2001) of about 472,000.

2.3.2 Physical Geography

This component describes the location and types of natural vegetative cover, aquatic habitats, and species habitats within the source protection area that are on the Species at Risk in Ontario List. It also describes the history, structure and composition of the surface, just below the surface, and deep beneath the surface (geology). In addition, this component describes natural landscape features (physiography), soil types, and surface shape and features (topography). Water movement on the surface (surface hydrology), such as rainfall, and water movement below the ground (groundwater hydrogeology), and climate, including air temperature and flooding are also included. A few details are given below but do not provide a complete picture of the characterization. For accurate descriptions, refer to the Thames Watershed and Region Watershed Characterization Report (2008).

Geology, Physiography and Soil Types

Bedrock is the rock formation deep under the ground, over which lies the overburden formation. The bedrock geology formations in Upper Thames River Source Protection Area are mainly the Dundee formation (fossiliferous limestone) and Detroit River Group (orthoquartzitic sandstone). The surficial geology is influenced by the type and nature of overburden. In the *UTRSPA*, the

Upper Thames River Source Protection Area Assessment Report

primary material of diamicton/till dominates, with silt plain north of St. Marys, and gravel near Komoka. Till is a mixture of clay, silt, sand and pebbles. The *UTRSPA* is mainly till plains without drumlins (streamlined landforms), with the exceptions of Oxford County which is a till plain with drumlins, and a sand plain north of London surrounded by spillways. As described below, till moraines are also important features of the *UTRSPA*. In the *UTRSPA*, 'silt & clay loam' type of soil is predominant (39%), with 'silt & clay' (26%) and 'loams' (15%) following. Maps 4, 5, 6 and 7 in Appendix 5 show the Thames watershed bedrock geology, overburden thickness, surficial geology and physiography respectively.

Topography, Hydrology and Hydrogeology

In the Upper Thames River Source Protection Area, the bedrock topography is higher than in other parts of the SPR, with the highest elevations occurring in northeastern parts of Perth County. Moraines are ridges of material that are generally topographic highs. Till moraines are seen across Komoka to Ingersoll and Woodstock, and in Perth County north of St. Marys, while a kame moraine is south of St. Marys. The Thames rises at three distinct points in the *UTRSPA*, near Mitchell (North Thames), Hickson (Middle Thames) and Tavistock (South Thames). The riverbeds are rocky and the valley slopes are steep, in contrast with the lower Thames River in the *LTVSPA* where the plains are flat. In the upper portion of the Thames River, the flow is 40% surface runoff and 60% 'base flow'. Base flow includes contributions from groundwater, tile drains, flow augmentation from reservoirs and treated sewage effluent discharge.

An aquifer is a water-bearing layer under the surface, which can be tapped by drilling groundwater wells. The Watershed Characterization Report Summary, included as Appendix 5 of this Assessment Report, includes a brief summary of the prevalent aquifers in the Upper Thames *SPA* while more detail is contained in the Watershed Characterization Report - Thames Watershed and Region (Thames-Sydenham and Region, December 2008). The Conceptual Water Budget (Appendix 6) includes a conceptualization of the aquifers and aquitards which continues to be improved through subsequent Water Budget work. Municipal groundwater systems in the Upper Thames *SPA* draw most of their drinking water from the bedrock aquifers with the exceptions of some Oxford and Middlesex systems which rely on water table aquifers. The aquifers tapped by each system are described in each system summary (Appendix 3). Similarly, private wells draw from both the water table and bedrock aquifers. In Perth County,

Upper Thames River Source Protection Area Assessment Report

wells draw predominantly from bedrock aquifers, while in Oxford and Middlesex, wells draw from both sources.

As with surface water, aquifers flow from an area of higher elevation (potential head) to an area of lower elevation. Maps 12 and 13 from the Watershed Characterization Report Summary (Appendix 5), illustrate the bedrock potentiometric surface elevation and the water table elevation across the region. The general flow direction in the Upper Thames *SPA* can be seen by referring to these maps. Groundwater flows generally from the higher areas in the northern end of the watershed towards the south and west parts of the region. Locally the aquifers, especially the water table aquifer, will flow towards lower hydrologic features such as streams. A more dramatic gradient is shown west of St. Marys where bedrock potentiometric surface drops off towards a karst area west of the watershed. Local aquifer flow directions can be seen by referring to the individual system *WHPA* maps 4-1-1 to 4-1-23 and considering that water flows from the outer edge of the *WHPA* towards the well as influenced by the local groundwater gradient and, to some degree, by the pumping of the wells.

Hydrology and climatic conditions are monitored locally by a combination of Environment Canada and Conservation Authority monitoring stations, including at London, Woodstock and Stratford in the Upper Thames River Source Protection Area. From plotting 10 year running averages over the data years of 1950 to 2005, an increasing level of precipitation in the 1970s and 1980s is seen with decreases recently. An increase in the linear trend line is seen at London, Woodstock and Stratford.

Natural Vegetative Cover

Wetlands are about 57 sq. km and make up less than 2% of the total *UTRSPA* watershed area of 3,447 sq. km, as shown in Map 23a of Appendix 5. Overall, wetland cover averages 1.7% with a high of 9.7%. The subwatersheds with the highest wetland cover are Black Creek (north of Stratford), Dorchester (east of London) and Komoka (west of London). All of these areas contain large wetland complexes. The vast majority of the remaining wetlands in the Upper Thames River Conservation Authority (UTRCA) are classified as deciduous swamps or mixed deciduous-coniferous swamps that are dominated by trees and shrubs such as silver maple, ash, willow, dogwood and cedar. Many swamps contain small pockets of marsh vegetation

Upper Thames River Source Protection Area Assessment Report

where emergent plants such as cattails, rushes and sedges dominate, but there are no large marsh sites. Bogs and fens are also very rare. There are a couple of kettle bogs in the London area. In the Upper Thames River watershed there are 31 provincially significant and 35 locally significant wetlands.

In the *UTRSPA* watershed, woodland/forest cover varies between 5 and 21% within the subwatersheds (with an average of 12%, or about 413 sq. km), as shown in Map 25a of Appendix 5. The subwatershed with the highest amount of woodland/forest cover is Dorchester, owing to the presence of the large Dorchester Swamp and North Dorchester Swamp complexes. The largest woodland/forest tract is Ellice Swamp (1,014 ha), located north of Stratford. The lowest amount of woodland/forest cover (4.9%) is in the North Mitchell watershed which is the headwaters of the North Branch of the Thames in Perth County.

The area of land adjacent to streams is often called the riparian zone or buffer zone. In the *UTRSPA*, the riparian areas ranged from a low of 6.1% to a high of 31.8% with an overall average of 21.14%. The lowest riparian woodland/forest cover occurs in the headwaters area of the North Branch of the Thames River in rural Perth County (North Mitchell and Whirl Creek subwatersheds). The highest cover is in the Dorchester watershed east of London and the River Bend and Oxbow Creek watersheds west of London.

Specific areas are protected from developmental changes that could alter their natural character. This protection is designated through federal, provincial and local initiatives. Depending on the degree of protection, “protected areas” are not likely to change over time and will encounter minimal human disturbance. There are several significant protected areas in the *UTRSPA* watershed. The Ellice Swamp covers approximately 856 hectares and is the largest woodlot in Perth County. Golspie Swamp covers 295 hectares and represents the third largest forested area remaining in Oxford County. The Dorchester Swamp is a 548-hectare site that is recognized as a Class 1 Significant Wetland, a Carolinian Canada Site and an Area of Natural and Scientific Interest (ANSI). The Sifton Bog in the City of London is a Class 2 provincially significant wetland and the most southerly large acidic bog in Canada. Also in London, Westminster Ponds/Pond Mills Conservation Area covers approximately 300 hectares with six major ponds over an area 3 kilometres long and 1.5 kilometres wide.

Upper Thames River Source Protection Area Assessment Report

Aquatic Ecology and Habitats

In the Thames Watershed & Region, the wide variety of habitats, favourable climate, nutrient-rich waters, and connection with the Great Lakes result in a particularly diverse aquatic community. The Thames River and its tributaries support one of the most diverse fish communities in Canada. Records exist for approximately 94 fish species in the Thames River subwatershed, which represents more than half of all of Ontario's 165 species. Table A5-1 (Appendix 5 Addendum) lists the fish species recorded in the Thames River subwatershed. Table A5-2 (Appendix 5 Addendum) lists the mussel species found in the Thames River.

Aquatic invertebrates, especially the benthic macroinvertebrates (BMI) that inhabit watercourse substrates, are abundant in all Thames reaches and tributaries. BMI communities consist of insect larvae, aquatic worms, crustaceans, and many other species. Most have fairly well known tolerances to pollution and disturbance. Table A5-3 (Appendix 5 Addendum) lists the benthic species commonly found in the Thames River and tributaries.

Introduced fish species found in the Thames such as the common carp and round goby are considered invasive species. In the Great Lakes, native freshwater mussel populations have been decimated by zebra mussels (from certain water bodies in Asia).

Map 20 of Appendix 5 illustrates the UTRCA Watershed Watercourse Classification to differentiate between municipal drains, natural watercourses (non-municipal drains) and some tiled (closed surface) watercourses. There are approximately 47% open municipal drains, 28% natural or non-municipal drains, and 25% tiled watercourses in the UTRCA watershed. The natural or non-municipal drains represent more than 25% of the length of watercourses in the UTRCA watershed. However, based on a review of Map 20 of Appendix 5, the majority of the natural watercourses are the main rivers including the Thames River, the north, middle and south branches of the Thames and the lower sections of some of the larger tributaries such as the Avon River. Approximately 24% of watercourses in this watershed provide suitable water quality and habitat conditions for sensitive species. Of those watercourses approximately 6% are municipal drains and 18% are natural watercourses.

Upper Thames River Source Protection Area Assessment Report

Table 2-2 shows the habitat and temperature assessment information for the *UTRSPA*. Some watercourses can be classified as both warmwater and coldwater, or as both permanent and intermittent flow. Sometimes thermal assessments were not possible, for example, during dry weather when there is no flow in certain watercourses. Approximately 10% of watercourses in *UTRSPA* are permanent cold/cool water streams with less than half considered to be natural. Approximately 61% are permanent warm water while about 30% are intermittent watercourses. Of the 61% that are warm water, there is an almost equal division between natural watercourses (31%) and municipal drains (30%). Of the roughly 30% of watercourses that are intermittent systems, or dry for most of the year, only 2% are considered natural while 28% are municipal drains.

Table 2-2 Habitat and Temperature Assessments in the UTRSPA

Municipality	Permanent Flow	Intermittent Flow	Coldwater	Warmwater
Biddulph	13	3	3	13
Blandford-Blenheim	7	27	0	31
Central Huron	2	0	2	0
East Zorra-Tavistock	44	50	18	78
Ingersoll	2	0	2	0
London	69	93	7	155
Middlesex Centre	71	111	6	176
Mitchell	6	0	0	6
North Dorchester	1	0	1	0
North Perth	0	8	0	8
Norwich	17	1	5	13
Perth East	82	79	18	143
Perth South	126	28	11	143
Sebringville	2	0	0	2
South Huron	16	3	0	19
South-West Oxford	25	18	9	34
St. Marys	13	1	0	14
Stratford	22	3	0	25
Strathroy-Caradoc	4	2	2	4
Thames Centre	84	120	20	184
West Perth	99	85	3	181
Zorra	115	145	43	217
Total	820	777	150	1,446

Much of the headwaters, particularly intermittent drains, have remnant pools that provide refuge areas for a variety of the more tolerant or hardy aquatic species. However changes such as the

Upper Thames River Source Protection Area Assessment Report

removal of cobble from the channels and the lack of pool riffles result in aquatic communities limited to hardy warmwater species.

In a few isolated headwater locations in the Thames, conditions allow more sensitive coldwater communities to persist. Sensitive BMI species, such as stoneflies and some caddisflies, are indicators of a high quality aquatic habitat or ecosystem. Coldwater fish species, such as trout and sculpin, that require well-oxygenated cool or cold flows year-round, can be found in these headwater streams. Moving downstream from the headwaters to medium-sized Thames tributaries, overall aquatic habitat generally improves as the stream size increases. In these streams, a diverse aquatic community is generally present, often including many mid-tolerant and the occasional sensitive BMI. In addition, most streams support a diverse fish community that may include top-level predators and Species at Risk (SAR). A few streams that have significant groundwater inputs support native brook trout or introduced brown trout populations.

Farther downstream, the larger tributaries and the three main Thames branches generally support aquatic communities of increased complexity and stability. Much of the north Thames, portions of the middle and south Thames, and a few larger tributaries support very diverse and productive aquatic communities. The communities of BMI are largely comprised of mid-tolerant and a few sensitive species. Top-level predators and species that require relatively clear flows and clean substrates to survive are well represented within the fish community. These river and stream reaches also provide habitat for a large proportion of the surviving fish, reptile and freshwater mussel SAR found in the Thames. From London downstream to the Delaware area, the flow and habitat conditions for the river are much like the upper branches but at a larger scale.

Impact of Human Activities on Aquatic Ecology and Habitats

The Watershed Characterization Report also discusses the impacts human activities have had on aquatic ecology and habitats. The Thames River is situated in a highly developed part of southern Ontario. The aquatic community faces many pressures from urban and rural land uses and human activities. Most of the watercourses have been greatly altered by human influences. On larger watercourses, many of the influences accrue from urban development, including channel alteration, bank hardening, storm water runoff, and sewage effluent input. Rural

Upper Thames River Source Protection Area Assessment Report

influences often involve smaller watercourses where habitat changes and alterations such as drains and channelization are aimed at improving agricultural operations. In general, species that prefer clear, fast flowing water are declining (Cudmore, B., C. A. MacKinnon and S. E. Madzia. Dec. 2004).

In 1986, Holm and Crossman completed a study comparing current (1985) information to historic surveys from the 1920s and 1940s. They identified water quality and fish habitat as conditions that had deteriorated significantly in the Thames River. They noted that turbidity and siltation had increased, and that stream flow rates had changed as a result of habitat disruptions such as impoundments. They also indicated a decline of species with a preference for clear, fast water and an increase in abundance of species more tolerant of turbidity.

Intermittent drain systems provide a significant function to the watershed. They provide fish habitat when wet and, in many cases, significant spawning areas during spring flooding. Some drains have pooled refuge areas (as evident in the Upper Thames watershed) and support habitat generalist species. These drains still support aquatic communities that primarily consist of tolerant BMI and fish species. These are particularly evident where agricultural best management practices (BMPs) have reduced agricultural impacts. In recent years, many of these intermittent watercourses have been converted to closed systems. The trend to close drain systems has altered the hydrograph, hydrologic regime and fluvial dynamics of the receiving watercourses and has led to an increase in erosion in downstream watercourses. Changes such as the removal of cobble from the channels and the lack of pool riffles result in aquatic communities limited to hardy warmwater species.

In a few isolated headwater locations in the Thames, more sensitive coldwater communities persist due to the presence of groundwater discharge, riparian vegetation and shading, headwater wetlands, and usually an undisturbed natural channel (although several drains support coldwater communities). Moving down from the headwaters, most medium-sized streams have natural channels or, if channelized, their stream power is often more in balance with the channel characteristics. Riffle/pool sequences have redeveloped with a firm (cobble/gravel/sand) substrate similar to that found in most natural watercourses. Most have an evident, well-defined flood plain with varying levels of disruption. A few have relatively

Upper Thames River Source Protection Area Assessment Report

undisturbed riparian vegetation and others are pastured or are idle pasture. The areas with idle pasture are now undergoing the slow process of natural succession or regeneration. Influences on these streams are silt, nutrient and pollutant inputs, both from neighbouring land use and from upstream sources. However, the improved habitat allows development of a complex and productive aquatic community with flood plain and hyporheic zone interactions. This enhances nutrient utilization and cycling. A diverse aquatic community is generally present. Farther downstream, the larger tributaries and the three main Thames branches generally support aquatic communities of increased complexity and stability. For the most part natural stream morphology and undisturbed flood plain is evident. The less impacted sections include much of the north Thames, portions of the middle and south Thames, and a few larger tributaries which support very diverse and productive aquatic communities. These river and stream reaches also provide habitat for a large proportion of the surviving fish, reptile and freshwater mussel Species at Risk (SAR) found in the Thames. However, many of these larger watercourses are influenced by urban development, including channel alteration, bank hardening, stormwater runoff, and sewage effluent input.

In general, a diverse community of mussels characterizes a healthy aquatic environment. There was once a diverse mussel community in the Thames. Mussel species that have disappeared, or mussel species that are extremely hard to find, indicate that aquatic conditions may be deteriorating. The primary threats to native freshwater mussel populations include turbidity, siltation, habitat loss or degradation, watercourse barriers, invasive species and poor water quality.

Species at Risk

A "species at risk" (SAR) is any naturally-occurring plant or animal in danger of extinction or of disappearing from the province (Source: <http://www.mnr.gov.on.ca/en/Business/Species/>). Map 29 of Appendix 5 shows the number and locations of SAR in the Thames watershed. Table A5-4 (Appendix 5 Addendum) lists the aquatic SAR (fish, reptile and mussel) in the Thames River subwatershed.

Of the more than 90 species of fish found in the Thames River watershed, there are currently 11 fish listed as Species at Risk (as of May 2010). The gravel chub is the only species considered

Upper Thames River Source Protection Area Assessment Report

to be extirpated (wildlife species no longer existing in the wild in Canada, but occurring elsewhere). Only two records exist for the Thames, and Canada, with the last specimen captured in 1958. A slight chance exists that the species may persist, as it is potentially very difficult to capture with the sampling methodology that has been utilized in recent years. The northern madtom is the only species listed as endangered (wildlife species facing imminent extirpation or extinction) by Committee on the Status of Endangered Wildlife in Canada (COSEWIC), although the eastern sand darter is listed as endangered on the provincial SAR list. Federally, the eastern sand darter, spotted gar, and black redhorse are threatened (wildlife species likely to become endangered if limiting factors are not reversed) species. The northern brook lamprey, grass pickerel, silver shiner, pugnose minnow, river redhorse, and spotted sucker are of special concern (wildlife species that may become a threatened or an endangered species because of a combination of biological characteristics and identified threats). Bigmouth buffalo, black buffalo and greenside darter were listed as special concern, but, have been delisted for a variety of reasons.

A few reptile SAR are identified in the Thames River watershed, amongst which the spiny softshell turtle (*Apalone spinifera spinifera*) was designated as threatened in 1989, by COSEWIC. The Thames River holds one of only three large communal nesting sites known in Canada, the other two occurring along the north shore of Lake Erie. The Thames is also one of the few rivers in Ontario from which consistent reports of the queen snake still occur. The queen snake was designated as endangered in 2010 by the Committee on the Status of Endangered Wildlife in Canada (COSEWIC).

There are ten SAR mussels (as of May 2010) in the Thames River: snuffbox, round hickorynut, kidneyshell, salamander mussel, rayed bean, rainbow, mapleleaf, wavyrayed lampmussel, fawnsfoot and round pigtoe. The snuffbox and round hickorynut are believed to be extirpated from the Thames River. Almost all of the mussels are listed as endangered by COSEWIC. The mapleleaf is threatened and the wavyrayed lampmussel has just recently changed from endangered to special concern. Provincially, under the Endangered Species Act for Ontario 2007, the mussels have a status of endangered, except for the rainbow and mapleleaf that are listed as threatened.

Upper Thames River Source Protection Area Assessment Report

2.3.3 Human Geography

The current population and estimated growth rate in each municipality are presented. The Upper Thames River Source Protection Area (*UTRSPA*) includes most of the municipality of City of London, and large parts of Elgin County, Middlesex County, Perth County and Oxford County. It also contains small parts of Huron County and Elgin County. The area covers approximately 3,423 square kilometres with a total watershed population (2001) of about 472,000. There are no First Nations in the *UTRSPA*, however there are First Nations in the other two SPAs of the source protection region.

Types of settlements (urban and rural centres) and land use (such as agricultural, residential and industrial) across the watershed are discussed. Map 1-4 of Appendix 1 shows the Areas of Settlement (as per the Places to Grow Act, 2005) in the Upper Thames River Source Protection Area. The largest urban centre within the Upper Thames River Source Protection Area is the City of London, with an approximate population of 336,539 in 2001. Table 2-3 shows the population by municipality, for the years 2006, 2001 and 1996. Map 2-1 in Appendix 1 shows the population density across the Upper Thames River watershed. Growth rate projections for municipalities in the *UTRSPA* are discussed in the Watershed Characterization Report.

Table 2-3 Population Density of Municipalities in the Upper Thames River SPA Watershed

Census Division	2006 Population	2001 Population	1996 Population	1996 to 2001 Population Change	2001 to 2006 Population Change
Middlesex (incl. City of London)	422,333	403,165	389,616	3.50%	4.70%
Elgin	85,351	81,553	79,159	3.00%	4.70%
Huron	59,325	59,701	60,220	-0.9%	-0.6%
Perth	74,344	73,675	72,106	2.2%	0.9%
Oxford	102,756	99,270	97,142	2.2%	3.5%

Map 30 of Appendix 5 shows the generalized land cover in the Thames Watershed & Region. Agriculture is the dominant land use, but a wide variety of industrial, commercial and institutional land uses also provide employment for people.

Upper Thames River Source Protection Area Assessment Report

General locations of federal lands in the Upper Thames River Source Protection Area are shown in Map 2-2 of Appendix 1. The Map 2-2 was generated using data available at the Treasury Board of Canada Secretariat website (<http://www.tbs-sct.gc.ca/dfrp-rbif/home-accueil.asp?Language=EN>), map navigator page.

Interaction between Physical and Human Geography

The watershed characterization report discusses the interaction between human and physical geography.

The original amount of wetland cover in the *UTRSPA* watershed is unknown. The total area of wetland cover (evaluated wetlands) is now less than 2% of the conservation authority's watershed area.

The smaller watercourses (first to third order streams) have, on average, 20% woodland riparian cover. The larger watercourses (fourth order and higher) have approximately 28% riparian cover. This difference is to be expected as many of the first and second order streams are farm drains that are often without any buffer at all.

Hardwood forests covered the majority of the upper Thames River watershed prior to European settlement with smaller pockets of grassland and savanna habitat. Species diversity was very high in this region due to the long growing season, rich soils and productive waterways. However, these same factors also made the area attractive for farming and urban development. Today, the woodland/forest cover in the *UTRSPA* is highly fragmented, existing as small woodlots separated by agricultural fields, urban development and other land uses. Over 70% of the woodlots in the Upper Thames watershed are less than 10 hectares. Due to the practice of clearing the acreage closest to the concession road for farming, many woodlots represent the 'back 40' of farms and are distributed in a linear fashion parallel to the roads.

Over the past century, several diseases and pests introduced by man have had a significant impact on the local tree species. The American Chestnut was destroyed by chestnut blight, caused by an Asian bark fungus accidentally introduced to America on imported Asiatic chestnut trees. The Blight was probably imported into North America from Asia in the early

Upper Thames River Source Protection Area Assessment Report

1900s. Similarly, the American Elm has been seriously affected by an introduced fungal disease, Dutch Elm Disease, with heavy mortality. The disease was accidentally introduced into North America in 1931, in shipments of logs from the Netherlands destined for use as veneer. More recently, the Emerald Ash Borer, which is native to China and eastern Asia, has left a path of destruction in Essex County in southwestern Ontario (as well as southeastern Michigan, northern Ohio and Indiana). It has been found in Chatham-Kent and Elgin County. Its significance for woodlands in the region is not yet known but ash trees form an important part of the local tree cover in many woodlots.

Oxford County is characterized by significant reserves of mineral aggregates from both naturally occurring sand and gravel deposits and bedrock-derived crushed stone. The presence of substantial, high-quality deposits has led to the establishment of significant quarrying and sand and gravel extraction industries in the county. Much of the activity is located in Zorra Township. Several sections of Middlesex County contain abundant Quaternary deposits of sand and gravel. Portions of the City of London, including the Byron area and land surrounding Fanshawe Conservation Area, contain valuable deposits in close proximity to a large market with high demands for aggregate resources. In the Komoka area of Middlesex Centre Township, most of the aggregate resources have been extracted, although some pits are still active. In Perth County, excessive overburden limits the accessibility of limestone deposits that underlay most of the county, with St. Marys being the notable exception. Here, extensive quarrying activity since the 1880s has produced a significant percentage of limestone used for the production of cement products in the Great Lakes region.

In the *UTRSPA* watershed, there are 11 active landfills, most of which are in Perth County. The active landfills are small and for local communities. In the *UTRSPA* watershed, there are brownfield areas in all of the major urban centres. Brownfields are generally abandoned or underused commercial or industrial areas thought to be contaminated by past activities but which have a potential for redevelopment. Municipalities have either developed or are developing policies and incentives to encourage brownfield re-development. In the smaller urban settings, brownfields are limited in size and location. The Brownfield and Community Improvement Plans (*CIP*) vary from municipality to municipality. In the City of London, the *CIP*

Upper Thames River Source Protection Area Assessment Report

for brownfields cover lands within the Urban Growth Boundary identified on the Land Use Map (Schedule “A”) of the Official Plan, allowing those lands possible remediation.

Southwestern Ontario has a long history related to the oil and gas industry. Map 31 of Appendix 5 shows the concentration of oil and gas wells across the area. Middlesex, Oxford and Perth Counties are underlain by Paleozoic sedimentary rocks that have the potential for occurrence of oil and gas resources. However, very few commercial discoveries of hydrocarbons have been made other than the Innerkip gas pool in the northwestern corner of the *UTRSPA* watershed. Natural gas has been produced from the Innerkip gas pool since 1961. Only non-commercial shows of oil or natural gas have been encountered in the rest of the watershed. Relatively few wells have been drilled over 100 m to explore for hydrocarbons in the area and there is potential for additional undiscovered pools.

Due to a number of factors, including moderate temperatures, adequate rainfall, adequate growing season and good soil, the major land use in the region is agricultural and, more specifically, cash crop land. As shown on Map 33 of Appendix 5, most of the soils in the region are Class 1, 2 or 3 soils that are suitable for the sustained production of common field crops. Over the last 40 years, a significant trend in the agriculture industry has been the conversion from a mixed land use of both livestock pasture and crop cultivation, to either of these. In the 20-year period from 1986 to 2006, the number of farm operations has decreased and the farm operation size has increased dramatically due to a number of factors. Farmland makes up over 80% of the land use in the region. Farming dates back more than 1,000 years in the Thames watershed, when the Woodland peoples grew corn on fertile flood plain lands. These areas were considered to be highly suitable for agriculture as well as trade, transportation and later, industry. In the *UTRSPA*, agriculture is the predominant land use in 26 of the 28 subwatershed units, and Oxford County is a highly productive area with several livestock and cash crop operations.

Navigation by vessels is limited to the Thames River below London. The Thames River, although not navigable for large craft, still provides a picturesque locale for recreational boating, canoeing, rowing, and kayaking. Canoeing can be done in much of the upper Thames and small boats can be used on the impoundments behind some of the dams on the river. Fishing takes

Upper Thames River Source Protection Area Assessment Report

place in these waters and many of the other creeks and streams in the region. The Wildwood, Pittock and Fanshawe dams provide flood control. Map 22 of Appendix 5 shows the dams and control structures in the Thames watershed. These structures may affect water quality and restrict fish migration. Along the shoreline of the Thames River are pockets of cottages as well as low density residential and estate residential development with direct access to both public and private roads. Permanent dwellings in these locations are attractive because of their scenic vistas, recreational amenities and relatively easy commute to urban centres.

There are 17 private and municipal/conservation authority campgrounds in the *UTRSPA*. There are more than 30 public and private golf courses located throughout the *UTRSPA* including several that incorporate lands (flood plains) adjacent to local watercourses.

As mentioned earlier, there are several significant protected areas in the *UTRSPA* watershed. Ellice Swamp in Perth County, Golspie Swamp in Oxford County, Dorchester Swamp in Dorchester and Middlesex County, Sifton Bog in the City of London, and the Westminster Ponds/Pond Mills Conservation Area in the City of London are some of the protected areas in the *UTRSPA*. These areas are protected from developmental changes that could alter their natural characteristics.

2.3.4 Water Quality

This component describes the water quality across the *SPA*. The selection of indicator substances (parameters) is discussed. The watershed inland surface water, the ambient groundwater, municipal well raw (untreated) water, and the municipal surface water intake raw water quality data is reviewed and assessed using certain standards or guidelines. Where possible, trend lines are shown and statistical analyses performed.

Both drinking water quality standards and aquatic protection guidelines are used to assess water quality. The comparison is only intended to provide a means of quality assessment by using an established value and is not intended to judge conformance of raw (untreated) water to the standards or guidelines. The operation of a water treatment plant, including treated and distributed water quality, is governed separately under the Safe Drinking Water Act.

Upper Thames River Source Protection Area Assessment Report

Inland Surface Water Quality Monitoring

To evaluate inland surface water quality, data from 24 surface water quality monitoring stations across the Upper Thames River Source Protection Area were reviewed for certain parameters. These stations are monitored under the Provincial Water Quality Monitoring Network (PWQMN) program. At all stations in the UTRCA subwatersheds, the current 75th percentile phosphorus levels are above the Interim Provincial Water Quality Objective (IPWQO) of 0.03 mg/L. Most sites are less than 0.15 mg/L but two stations, Thames Woodstock and Thames Byron, are over 0.2 mg/L. The historic phosphorus values are generally between two and eight times the IPWQO. Seven of 30 UTRCA stations have current 75th percentile nitrate values that are higher than the Ontario treated drinking water standard (ODWS) of 10 mg/L. To evaluate water quality for aquatic life, a nitrate level of 2.93 mg/L has been used for comparison. Only two UTRCA stations have current 75th percentile nitrate values that are less than 2.93 mg/L. Historically, the levels at all UTRCA stations were above 2.93 mg/L with the exception of Dorchester Swamp Creek which has been consistently below 2.93 mg/L.

The current 75th percentile chloride concentrations at all UTRCA sites are below the ODWS of 250 mg/L. Historically, most UTRCA sites have concentrations that are below 100 mg/L. However, the chloride concentrations at the Avon River have shown a significant increase since 1990-1994 from approximately 100 mg/L to over 200 mg/L. The Environment Canada aquatic health guideline for chloride is 210 mg/L. The current 75th percentile chloride concentration at the Avon River station in the UTRCA watershed is close to this and 25% of the samples are above 210 mg/L. All other UTRCA stations have chloride concentrations below 210 mg/L.

A review of data for *Escherichia coli* (*E. coli*), a bacterial indicator, shows that the indicator is consistently above the recreational use guideline of 100 counts per 100 mL at most of the monitoring stations. The recreational use guideline for *E. coli* is intended for application by the local Medical Officer of Health, to decide the suitability of the use of swimming and bathing beaches. Copper and zinc levels are lower than their respective guidelines at all stations. At the UTRCA stations in 1980-84, all stations had samples above the lead ODWS of 10 µg/L and the percentage of samples above the ODWS ranged from 21% to 51%. By 2000-04, five of the six stations with historic data had all sample results below the ODWS and the other station only had 2% of the samples above the ODWS.

Upper Thames River Source Protection Area Assessment Report

Groundwater Quality Monitoring

To evaluate the groundwater quality, data from 23 groundwater monitoring wells across the Upper Thames River Source Protection Area were reviewed. The monitoring is part of the Provincial Groundwater Monitoring Network (PGMN) program. Based on the PGMN monitoring, the UTRCA overburden and bedrock wells are dominated by calcium-magnesium-bicarbonate water. The carbonate, magnesium (and sulphate) ions in the groundwater primarily originate from the carbonate rock at depth (bedrock: dolostones, limestones and evaporites) and the carbonate material incorporated in overburden sediments.

Fluoride levels were above the ODWS of 1.5 mg/L in seven UTRCA monitoring wells. Fluoride can occur naturally in groundwater at levels that are above the drinking water standard. Two monitoring wells had nitrate + nitrite or nitrate results above the ODWS of 10 mg/L, one monitoring well had arsenic above the ODWS of 0.025 mg/L, and one monitoring well had cadmium above the ODWS of 0.005 mg/L. In the UTRCA, 13 monitoring wells had iron above the Ontario Drinking Water aesthetic objective (AO) of 0.3 mg/L and six monitoring wells had manganese above 0.05 mg/L. Iron and manganese in groundwater are usually due to the natural weathering of rocks and minerals. In the UTRCA watershed, dissolved solids (five monitoring wells), dissolved organic carbon (three monitoring wells) and chloride (one monitoring well) were found above the AOs. Sodium values above 20 mg/L can be of concern for individuals that are on a sodium restricted diet and the local Medical Officer of Health is notified. Sodium above 20 mg/L (but below the ODWS of 200 mg/L) was found in nine UTRCA monitoring wells. There were 19 UTRCA monitoring wells that had hardness levels that were not within the Ontario Drinking Water Operational Guideline (OG) range of 80 to 100 mg/L. In addition to hardness, four UTRCA monitoring wells had high aluminum and one UTRCA monitoring well had high pH.

Municipal Groundwater System Water Quality

In the Upper Thames River Source Protection Area, there are 22 municipal groundwater and no surface water drinking water systems. Data used to evaluate water quality of raw water to the drinking water systems were: Drinking Water Surveillance Program (*DWSP*), Drinking Water Information System (*DWIS*), Annual Drinking Water System Reports, Ministry of Environment Inspection reports and minimal water treatment plant laboratory data. Similar to the findings of the groundwater monitoring data review, fluoride in raw (untreated) municipal well water is

Upper Thames River Source Protection Area Assessment Report

generally higher than the treated drinking water standard. Bacterial indicator total coliform is present in untreated water of most municipal wells, but *E. coli* was present in untreated water of only five systems. Raw (untreated) water quality is compared against treated drinking water standards and this comparison does not reflect the treated drinking water quality. The treated drinking water standards include those from the Reg. 169/03, and Table 4 parameters from the Technical Support Document for the Ontario Drinking Water Standards, Objectives and Guidelines (MOE 2006).

Nitrates occur above the Reg. 169/03 treated drinking water standard of 10 mg/L in the Thornton wellfield of the Woodstock wells. Fluoride occurs above the Reg. 169/03 treated drinking water standard of 1.5 mg/L in most wells, and is known to be naturally occurring in the groundwater.

In Middlesex County and City of London back up wells, sodium was below the Table 4 treated drinking water standard of 200 mg/L but above the 20 mg/L Health Unit notification level. One instance of fluoride above Reg. 169/03 treated drinking water standard of 1.5 mg/L occurred in the Thorndale system in 2004. In Oxford County, the Embro, Thamesford, Woodstock and Ingersoll systems had wells that had sodium levels below the Table 4 drinking water standard of 200 mg/L but over the Health Unit notification level of 20 mg/L. The Ingersoll, Lakeside and Thamesford systems had wells that had instances of fluoride above the Reg. 169/03 treated drinking water standard.

In Perth County, St. Marys and City of Stratford systems, the Sebringville, St. Pauls and Stratford well systems, sodium levels above the 20 mg/L Health Unit notification level are observed but the sodium levels are below the Table 4 drinking water standard of 200 mg/L. The Mitchell, Sebringville, St. Pauls and Stratford well systems had instances of fluoride levels above the Reg. 169/03 treated drinking water standard. In the Sebringville and Stratford systems, fluoride levels are noticeably high, often above 2.0 mg/L.

In addition, in Middlesex County, iron is above the Table 4 treated drinking water standard in the Kilworth-Komoka (now decommissioned) and Melrose systems, with manganese above Table 4 treated drinking water standard also in the former. Iron and manganese levels above the ODWS are also observed in a few Oxford County wells. In the City of London back up wells, organic

nitrogen and phosphorus also are above Table 4 treated drinking water standard and IPWQO, respectively. Hardness and total dissolved solids are seen in City of London and St. Marys wells. In Perth, iron levels above the Table 4 treated drinking water standard are observed.

2.3.5 Water Quantity

In this component, the water use across the Source Protection Area is discussed. Section 34 of the Ontario Water Resources Act (*OWRA*) requires anyone taking more than a total of 50,000 litres of water per day to acquire a Permit To Take Water (*PTTW*). Water taking includes uses which return the water to the source, as well as those which do not. Water taking also includes taking water into storage. In Section 3 - Water Budget and Water Quantity Stress Assessment water use is considered in more detail, including quantifying how much of the water taking is consumptive.

Water takers have a responsibility to ensure that the amount of water they use does not threaten the environment or existing water users. Some water takings are exempt from the requirement to obtain a permit. These include takings by an individual for ordinary household purposes, and water takings for the direct watering of livestock or poultry or for firefighting purposes. The approximate water taking (use) by sector (agricultural, commercial, industrial, municipal, water supply, dewatering, remediation, construction) is presented and described. The water taking for each subwatershed catchment area is also presented. These catchment areas are delineated through the Conceptual Water Budget study, which is described in Section 3 - Water Budget and Water Quantity Stress Assessment.

In the Thames watershed and region, while the agricultural sector has around 33% of the total permits, the percent of total maximum volume permitted is only 5%. This difference probably reflects the seasonal nature of the water taking associated with crop irrigation. Water supply makes up about 24% of the water taking permits, and includes takings by municipalities, campgrounds and communal uses.

2.3.6 Drinking Water Systems

There are 23 municipal drinking water systems which service people living in the Upper Thames River *SPA* of which two are located outside the *SPA*. The drinking water supply systems

Upper Thames River Source Protection Area Assessment Report

servicing the Upper Thames River *SPA* are shown in Map 1-3 of Appendix 1. Details are provided in Table 2-4. The 21 municipal drinking water systems located within this *SPA* are included in the Upper Thames River Source Protection Area Terms of Reference document, and therefore in this Assessment Report.

As per Regulation 287/07 (Section 14), an existing or planned major residential system may be exempted from Section 15 (2) (e) (i) of the Clean Water Act (i.e. identifying wellhead protection areas and intake protection zones), if the council of the municipality that owns the system has:

- passed a resolution stating that the municipality intends, within five years after the day the resolution is passed to discontinue the use of the drinking water system, and to make an application under the Safe Drinking Water Act, 2002 (and the Ontario Water Resources Act, if an IPZ or WHPA is delineated) for the revocation of any approval, municipal drinking water licence or drinking water works permit that is applicable to the drinking water system;
- published notice of the resolution referred to in clause (a) in one or more newspapers that, in the opinion of the council of the municipality, are of sufficiently general circulation to bring the notice to the attention of the public in the municipality; and
- sent a copy of the resolution referred to in clause (a) to the source protection committee for the source protection area

In June 2009, the Municipality of Strathroy-Caradoc passed a council resolution in order to exempt the Mt Brydges groundwater drinking water system from consideration in further technical studies towards the Assessment Report. Water is now obtained from the Lake Huron Primary Water Supply.

The Kilworth-Komoka wells were decommissioned in October 2010. The community of Komoka/Kilworth is now supplied by the Lake Huron Primary Water Supply System.

Upper Thames River Source Protection Area Assessment Report

Drinking Water System (No. of wells)	System Type*	Operating Authority	Approx Population Served	Well Name	Well Depth (m)	Pumping Rates m ³ /day		
						Max. Annual	Avg. Annual	Avg. Monthly
Birr (1)	2	American Water Systems	63	Well #1	49	4743	4316	4316
Dorchester (9)	1	Municipality of Thames Centre	5586	Well 2PW-1	18	152624	124982	10415
				Well 3PW-1	10	90684	76351	6363
				Well 3PW-2B	12	115904	100932	8411
				Well 3PW-3	12	104974	86163	7180
				Well 3PW-4A	12	119575	94716	7893
				Well 3PW-7	12	100963	67518	5626
				Well 3PW-5	27	421	335	23
				Well 3PW-6	27	1003	688	48
				Well 3PW-8 (New well)	13.1	NA	71508	5959
London (7)	1	Corporation of the City of London	Back up	Fanshawe 1 (Standby)	13.4	NA	1671335	139278
				Fanshawe 2 (Standby)	12.6	NA	1671335	139278
				Fanshawe 3 (Standby)	14.8	NA	1671335	139278
				Fanshawe 4 (Standby)	16.1	NA	1671335	139278
				Fanshawe 5 (Standby)	12.2	NA	1671335	139278
				Fanshawe 6 (Standby)	12.2	NA	1671335	139278
				Hyde Park Well (Standby)	37.3	NA	1671335	139278
Melrose (2)	2	American Water Systems	224	Well 2	25.6	17263	12248	1017
				Well 3	24.7	13882	11931	994
Thorndale (2)	1	Municipality of Thames Centre	675	Well 1	46	23891	20026	1669
				Well 2	45.1	23338	19376	1615

Upper Thames River Source Protection Area Assessment Report

Drinking Water System (No. of wells)	System Type*	Operating Authority	Approx Population Served	Well Name	Well Depth (m)	Pumping Rates m ³ /day		
						Max. Annual	Avg. Annual	Avg. Monthly
Beachville (1)	2	County of Oxford	180	Well #1	91.5	24156	21092	1758
Embro (2)	1	County of Oxford	828	Well 1	60	46248	38049	3171
				Well 3	60	46693	43282	3607
Hickson (1)	1	County of Oxford	99	Well # 2	53	12087	11111	926
Ingersoll (7)	1	County of Oxford	13,572	Well 2	140.5	549028	478392	39866
				Well 3 (Not presently in use)	117	259407	141017	15998
				Well 5	108.9	559446	493549	42714
				Well 7	122.8	188504	66668	7757
				Well 8	125.3	638316	378813	31568
				Well 10	112.5	1174171	909965	76935
				Well 11 (offline)	115.7	NA	NA	NA
Innerkip (2)	1	County of Oxford	935	Well 1	35	46938	41828	3486
				Well 2	35	47604	44285	3690
Lakeside (1)	1	County of Oxford	310	Well 2	106	16620	15851	1321
Mt Elgin (2, includes 1 planned well)	1	County of Oxford	369	Well #3	60	36536	34757	2933
				Graydon well (planned)	No Data	No Data	No Data	No data
Tavistock (3)	1	County of Oxford	2658	Well 1	19.5	30873	24059	2005
				Well 2A	48	171953	62004	5167
				Well 3	61.5	447789	350055	34510
Thamesford (3)	1	County of Oxford	2016	Well #1	9.4	203581	105480	8790
				Well #2	14	203581	81069	11260
				Well #3	78	203581	132310	11026
Woodstock (11, includes	1	County of Oxford	36,600	Well #1 (Thornton)	30	1980288	767680	63973

Upper Thames River Source Protection Area Assessment Report

Drinking Water System (No. of wells)	System Type*	Operating Authority	Approx Population Served	Well Name	Well Depth (m)	Pumping Rates m ³ /day		
						Max. Annual	Avg. Annual	Avg. Monthly
1 planned well)				Well #2 (Tabor)	20.8	1945843	1382113	115176
				Well #3 (Thornton)	16.1	275460	134961	11247
				Well #4 (Tabor)	23.5	2148179	1627792	135649
				Well #5 (Thornton)	27.1	1666172	444085	37007
				Well #6	48.8	609105	300616	25051
				Well #7 (Offline)	62.5	5492	5492	458
				Well #8 (Thornton)	14.6	776768	542950	45246
				Well #9	63.1	8581	5841	487
				Well #11 (Thornton)	31.9	1089339	791825	65985
				Bond well (planned)	No Data	No Data	No Data	No data
Mitchell (4)	1	West Perth Power Inc.	4000	Well 1 (Standby)	24.4	NA	69350	5779
				Well 2 (Standby)	33.5	NA	69350	5779
				Well 3	60	204458	989915	32493
				Well 4	56	567419	801726	66811
Sebringville (1)	2	Township of Perth South	90	Well 1	54.9	15152	12442	1037
Shakespeare (1)	2	Township of Perth East (AWS)	220	Well 1	85	1158	851	71
St. Marys (3)	1	Town of St. Marys	6200	Well 1	45.7	695162	615372	51281
				Well #2A	44.5	238348	214526	17877
				Well I#3	44.5	680292	431474	35956
St. Pauls (1)	2	Township of Perth South	90	Well #1	70.4	8389	8147	679
Stratford (11)	1	City of Stratford	30,460	Well (Chestnut)	150.3	83650	68221	5685

Upper Thames River Source Protection Area Assessment Report

Table 2-4 Municipal Drinking Water Systems Serving the Upper Thames River Source Protection Area

Drinking Water System (No. of wells)	System Type*	Operating Authority	Approx Population Served	Well Name	Well Depth (m)	Pumping Rates m ³ /day		
						Max. Annual	Avg. Annual	Avg. Monthly
				Well (Dunn)	135.3	1145360	996574	83048
				Well (Lorne)	137.5	198005	116183	9682
				Well (Mornington)	99.4	418440	205663	17139
				Well (O'Loane)	135.3	798890	661318	55110
				Well 1 (Romeo)	122	357747	280073	23339
				Well 2 (Romeo)	121	239425	137050	11421
				Well 3 (Romeo)	107	299028	186700	15558
				Well 4 (Romeo)	103	117393	85855	7155
				Well 6 (Romeo)	139	1289283	1080007	90001
				Well 7 (Romeo)	81	746575	574866	47905
Lake Huron Primary Water Supply System**								
Elgin Area Water Supply System***								

*System Type 1 – Large municipal residential, 2 – small municipal residential

**Serve some area of the UTRSPA but located outside of this SPA and therefore not included in this Assessment Report; see the Ausable Bayfield Source Protection Area Assessment Report for information

*** Serve some area of the UTRSPA but located outside of this SPA and therefore not included in this Assessment Report; see the Kettle Creek Source Protection Area Assessment Report for information

Residents in the northern part of the Thames watershed rely on treated groundwater for their drinking water. These include the communities of Shakespeare, Mitchell, St. Pauls and Sebringville in Perth County, City of Stratford and Town of St. Marys, and communities of Beachville, Embro, Hickson (King subdivision), Ingersoll, Innerkip, Lakeside, Mt Elgin, Tavistock, Thamesford, Woodstock and Sweaburg in Oxford County.

Upper Thames River Source Protection Area Assessment Report

Some parts of Middlesex County (communities of Birr, Melrose in Middlesex Centre, and Thorndale and Dorchester in Thames Centre) also have municipal systems that use groundwater sources.

Residents in the City of London and some neighbouring Middlesex communities (including Delaware and Ballymote) use treated surface water piped from Lake Huron (Lake Huron Primary Water Supply System) and Lake Erie (Elgin Area Water Supply System). The Fanshawe and Hyde Park well systems serve as back up drinking water systems to the City of London. Most of the water for residents in Elgin County is from Lake Erie.

2.4 Data Gaps

The data gaps encountered during the preparation of the watershed characterization report are listed in Table 2-5 below.

Upper Thames River Source Protection Area Assessment Report

Table 2-5 Watershed Characterization Data Gaps relevant to the Upper Thames River Source Protection Area

Subject	Data Gaps
Aquatic Ecology	
Fisheries Evaluation	Cold water refuges in natural water systems, historic evidence of cold water streams has not been investigated, application of indices such as the Index of Biological Integrity (IBI) to existing fish data.
Aquatic Macroinvertebrates - Habitat Conditions & Water Quality	Simpson's Diversity Index should also be considered, Analysis of physiography & land use to identify potential communities and groundwater quality/quantity stressors and impacts.
Reptile - Survival habitat and population dynamics	Extent, abundance and population demographics of prey (needed for some species); lack of species information, habitat identification, seasonal dispersal, population isolation, reproductive success, past distribution.
Species at Risk - Range and numbers of fish species at risk	Sections of the Thames River have little or no sampling, population, abundance, distribution or status unknown for some species.
Human Characterization	
Landfills	Information on active, closed landfills and expansions not provided for the Upper Thames watershed.
Water Quality	
Inland surface water quality - physical, chemical and microbial	Data from 1997 to 2001 from the commonly used provincial water quality monitoring network (PWQMN) dataset does not exist.
Additional sources of information	COA, Health Unit, sediment analysis and Research data have not been reviewed.
Inland and intakes surface water and groundwater quality – emerging pollutants	Not enough data on emerging contaminants (fire retardants, pharmaceuticals, algae toxins, etc.).
Groundwater monitoring well data	Comprehensive data not available.
Long term municipal groundwater well physical and chemical data	Comprehensive, long-term data not available; alternate sources of data used.
Wildlife impact on water quality	Locations of large populations of wildlife and the resulting effect on water quality (pathogen contamination and nutrient loading) require a better understanding.
Water Quantity	
Permit To Take Water Data	Data out of date - Many permits in database have expired dates and it is unclear if they have been renewed.
Water uses	Data Incomplete - Older permits only have maximum water taking per day. Difficult to determine actual usage.

Section 9.0 Data Gaps and Next Steps lists those data gaps considered to be a priority in filling, in order to meet the requirements of the Assessment Report. From the above table, the 'water uses' data gap is brought forward to Section 9.0.